

Marking behavior of Asiatic black bears at rub trees

Yoh Ogawa^{1,6}, Kahoko Tochigi²,
Tomoko Naganuma³, Bainah S. Dewi⁴, and
Shinsuke Koike^{3,5}

¹Tokyo Metropolitan Agricultural High School, 1-10-2
Kotobuki, Fuchu 183-0056, Japan

²United Graduate School of Agriculture, Tokyo
University of Agriculture and Technology, 3-5-8 Saiwai,
Fuchu 183-8509, Japan

³Institute of Global Innovation Research, Tokyo
University of Agriculture and Technology, 3-5-8 Saiwai,
Fuchu 183-8509, Japan

⁴Tropical Biodiversity Research and Development
Center, University of Lampung, Bandar Lampung 35145,
Indonesia

⁵Institute of Agriculture, Tokyo University of Agriculture
and Technology, 3-5-8 Saiwai, Fuchu 183-8509, Japan

Abstract: We studied the characteristics of the marking behavior of Asiatic black bears (*Ursus thibetanus*) at rub trees. We recorded the tree-marking (tree-rubbing and bark-scratching) and associated (stomping) behaviors in Japan between 2010 and 2015 using automatic cameras and compared the results with those from previous studies on other bear species. We recorded 22 visits to trees by ≥ 6 different individuals. During these visits, there were no differences in the frequency or duration between the 3 body orientations of back rubbing, front rubbing, and body-side rubbing, which differs from the findings for other bear species. However, bipedal standing was the most common posture, as observed in other species. We also analyzed the order of tree-marking actions. For a better understanding of the Asiatic black bear's marking behavior, further examples of marking behavior and anatomical studies of secretory glands are needed.

Key words: Asiatic black bear, body orientation, Japan, marking, scent-marking, stomping, tree-rubbing, *Ursus thibetanus*

DOI: 10.2192/URSUS-D-20-00028.1
Ursus 32:article e24 (2021)

Among the Ursidae, tree-rubbing has been mainly studied in brown bears (*Ursus arctos*; e.g., Green and Mattson 2003, Clapham et al. 2012, Sato et al. 2014, Tat-

toni et al. 2015), American black bears (*U. americanus*; e.g., Burst and Pelton 1983, Taylor et al. 2015), Andean bears (*Tremarctos ornatus*; e.g., Filipczyková et al. 2016, Kleiner et al. 2018), and giant pandas (*Ailuropoda melanoleuca*; e.g., White et al. 2002, Nie et al. 2012). Tree-rubbing behavior in bears involves multiple individuals repeatedly rubbing the same tree, sometimes accompanied by bark-scratching with claws and teeth and urinating (Karamanlidis et al. 2007, Kendall et al. 2009, Taylor et al. 2015, Filipczyková et al. 2016).

Brown bears and American black bears rub parts of their bodies against artificial objects as well, such as creosoted power poles and fence posts (e.g., Karamanlidis et al. 2007). The function of bears' rubbing behavior is not yet fully understood, and there are a variety of speculations. One category is physiological drives experienced by the bears, which includes reasons such as reactions to skin parasites, inflammation, itching, or molting, as well as physical stimulation or curiosity (e.g., Green and Mattson 2003, Taylor et al. 2015). Another category relates to tree-marking with animal scents and is based on the observations that multiple bears may rub the same tree, that males are more likely to rub trees than females, and that tree-rubbing peaks during the breeding season (e.g., Green and Mattson 2003, Clapham et al. 2012, Taylor et al. 2015). Some researchers believe scent-marking is used for communication for these solitary animals, which are active across large areas (Clapham et al. 2012, 2014; Sato et al. 2014).

Recent detailed behavioral and physiological analyses of brown bears have indicated that differences in their actions and postures during tree-rubbing may be related to differences in scent-marking by glandular secretions. For example, Clapham et al. concluded that back rubbing (BR) on 2 legs was the core marking posture, based on observations of the motor activity of tree-rubbing, including analysis of the order of behavior with high probability. In addition, oily secretions and enlarged sebaceous glands have been found on the skin of the back during the breeding season, which suggests that the sebaceous glands on the back may be involved in BR (Tomiyasu et al. 2017). Furthermore, bears have been observed smelling the ground before tree-rubbing and sitting in front of the tree after tree-rubbing, and young males are able to distinguish mature males by their anal secretions (Rosell et al. 2011, Jojola et al. 2012). Also, mature males engage in pedal marking behavior, whereby a bear walks with a stomp and leaves distinctly recessed footprints (Green

⁶email: onwogawa@mac.com

and Mattson 2003; Clapham et al. 2013, 2014; Sato et al. 2014), before and after tree-rubbing and other marking behavior.

In recent years, it has become clear that Asiatic black bears (*Ursus thibetanus*) also perform tree-rubbing (Latham et al. 2012). Indeed, multiple bears have been recorded rubbing against the same conifer tree (Ogawa et al. 2020), but there has been no detailed analysis of Asiatic black bear rubbing behavior to date, and there are no reports of studies on secretory glands.

The aim of this study was to study the characteristics of tree-marking behaviors in Asiatic black bears and compare them with those of other bear species. Previous studies in brown bears have shown that the order in which tree-marking behavior occurs varies by sex or age of the bear (Clapham et al. 2014) and may also vary among different species of bears. Thus, we focused on the characteristics of body-rubbing and bark-scratching behaviors (body posture, body orientation, and order of actions). We then compared the characteristics of tree-marking behaviors in Asiatic black bears and other bear species as an important step toward understanding the role of this behavior in intraspecific communication.

Study area

We conducted this study in Okutama (35°48'N, 139°5'E), the westernmost suburb of Tokyo, Japan (Fig. S1, Supplemental material). The region has a Pacific Ocean-type climate, with heavy rainfall in summer and little snow in winter. The mean annual precipitation is 1,510 mm, and the mean annual temperature is 12.4°C, ranging from 0.6°C in January to 24.2°C in August (2006–2017; Japan Meteorological Agency 2018).

The study area is mainly covered with natural forests (40%) and conifer plantations (50%; Koike et al. 2008). The natural forests are dominated by *Castanea crenata* and *Quercus serrata* in the lower mountain zone (400–500 m above sea level [a.s.l.]); *Q. crispula*, *C. crenata*, and *Fagus crenata* in the middle zone (500–1,500 m a.s.l.); and *Abies homolepis* and *Tsuga diversifolia* in the upper zone (1,500–1,800 m a.s.l.). The conifer plantations include Japanese cedar (*Cryptomeria japonica*), Japanese cypress (*Chamaecyparis obtusa*), and Japanese larch (*Larix kaempferi*). We conducted the present camera survey in a mixed plantation of Japanese larch and Japanese cedar (elevation range: 950–1,000 m a.s.l., area size: 20,000 m²).

Materials and methods

Camera setup

To record the tree-marking behavior of bears in the study area, we installed automatic video cameras (HCO Scoutguard SG550 [HCO Outdoor Products, LLC., Johns Creek Pkwy, Georgia] and Bushnell HD Max [Bushnell Corporation, Overland Park, Kansas]) and photo cameras (Reconyx RC55 [Reconyx, Inc., Holmen, Wisconsin]) pointing at Japanese larch and Japanese cedar trees that had bark damaged from antler sharpening by sika deer (*Cervus nippon*) and on which bear hairs had previously been found (unpublished data). We conducted the survey on 16 different trees between 2010 and 2015: 5 Japanese cedar in 2010; 4 Japanese cedar and 2 Japanese larch in 2011; 2 Japanese larch in 2013; and 2 Japanese larch and 1 Japanese cedar in 2015. The cameras were generally in place between early May and early November in the survey years and were installed 2–5 m away from the target trees at a height of 1–2 m above the ground. We installed photo cameras on 3 Japanese cedar and 1 larch in 2011, and installed video cameras on all the rest. We set the video cameras to take 60-second videos each time they were activated, with 5-second intervals between recordings; whereas, we set the photo cameras to take a series of 10 still pictures at 1-second intervals as a single camera activation, with 5-second intervals between series. During each survey period, we changed memory cards and batteries once per month.

Classification of actions

In the video recordings and captured images, we defined any behavior in which part of an individual bear's body touched a target tree (specifically, tree-rubbing or bark-scratching) as tree-marking, and we regarded any tree-marking behavior that was captured during a single camera activation as one visit. In cases where 2 individuals performed simultaneous tree-marking, we considered that as 2 visits. If tree-marking by the same individual was captured within 30 minutes of the end of a preceding video sequence, we considered it as a single visit. We identified gender by external genitalia, and identified individuals on the basis of a number of observed details, including chest markings, degree of body hair depletion, and physique, according to the Individual Identification Reporting Checklist (Choo et al. 2020).

We classified the characteristics of the tree markings as shown in Figure S2 (Supplemental material) and recorded the number of times each action was performed for each visit. We classified “tree-rubbing” as a type of tree-marking behavior that involved 1 of the following

5 actions, which were based on a combination of posture (which relates to the height on the tree where the body part was rubbed) and body orientation (which relates to the part of the body that is rubbed): rubbing the side of the body and neck against a tree with all 4 feet in contact with the ground (quadrupedal standing position, body-side rubbing [4BsR]); rubbing the front of the body against a tree with 2 feet in contact with the ground (bipedal standing position, front rubbing [2FR]); rubbing the back of the body against a tree with 2 feet in contact with the ground (bipedal standing position, back rubbing [2BR]); sitting while rubbing the front against a tree (sitting position, front rubbing [SFR]); and sitting while rubbing the back of the body against a tree (sitting position, back rubbing [SBR]). We classified “bark-scratching” as another type of tree-marking behavior that comprised 2 actions: scratching bark while standing on 2 feet (bipedal standing position, bark-scratching [2Sc]) or scratching bark while sitting (sitting position, bark-scratching [SSc]). We included stomping (St) as an associated behavior in our observations, whereby the bear, after leaving the tree, walked heavily around over the same spot with both the front and back feet while looking down and smelling for ground odors.

We used Quick Time Player 7 (Apple Inc. Version 7.6.6, Cupertino, California) to analyze the duration, frequency, and order of each of the aforementioned 8 actions recorded during each visit. We treated the same action observed to continue across camera activations as a single action, whereas we treated the same action performed again after other actions as a separate action.

Data analysis

We conducted multichannel attribution analysis to investigate whether observable patterns of behavior occurred during a visit to a target tree. This type of analysis determines the starting point of the behavior, the specific intermediate actions performed during the behavior, and the end point of the behavior in order of action, and also divides the sequence from the start point to the end point into 2 points and calculates the probability that these sequences of events will be passed through. We used the Markov chain model to investigate the probability that a bear would pass through each action sequence, whereby we placed the starting point before the behavior labeled “tree-marking,” we labeled the 7 intermediate actions, and we set “St” or “leaving without St” as the end point. We constructed the model using the Channel Attribution package (Altomare and Loris 2018) and displayed the re-

sults using the visNetwork package (Thieurmél 2016) in Program R (R Core Team 2019).

We analyzed differences in the frequency of occurrence of and mean percentage of time spent in (hereafter, “percentage duration” of) each body orientation (namely FR [front rubbing], BR [back rubbing], or BsR [body-side rubbing]) and posture (namely 2R [bipedal standing position, rubbing], 4R [quadrupedal standing position, rubbing], or SR [sitting position, rubbing]) using 2×2 chi-squared tests. Specifically, these tests compared the differences in the percentage of a visit in which a given bear displayed FR versus BR, FR versus BsR, BR versus BsR, 2R versus 4R, 2R versus SR, and 4R versus SR. In addition, we compared differences in the dispersion of the percentages of time spent in each body orientation or posture during the visits in which each behavior was observed using a *t*-test.

Results

With the automatic cameras, we observed ≥ 6 different individual bears visiting 5 of the 16 target trees (2 Japanese cedar and 3 Japanese larch) 22 times between May and October across all years. There were 13 visits from males, 1 visit from a female accompanied by 1 offspring (gender unknown; counted as 2 visits), and 7 visits from bears of undetermined gender.

We observed 74 instances of tree-marking behavior over the 22 visits. The average tree-marking rate was 3.4 ± 2.9 times/visit (range = 1–12 times). The average tree-marking duration per bear was 33.8 ± 37.9 seconds (1–161 sec), and there was a positive correlation between the number of times tree-marking was performed per visit and the visit duration ($r = 0.96$). Not all visits resulted in established tree-marking behaviors (Table 1). The analysis of body orientation revealed that there was no difference in the percentage of visits in which BR, FR, and BsR were displayed (59.1% each). By contrast, the analysis of body postures showed that 2R was displayed in 72.7% of visits, significantly more frequently than 4R (59.1%; $\chi^2 = 4.14$, $P = 0.042$) and SR (40.9%; $\chi^2 = 20.6$, $P < 0.001$), and 4R was displayed significantly more frequently than SR ($\chi^2 = 6.61$, $P = 0.010$). In terms of body orientation and posture, there was no significant difference in the percentage duration between FR and BR or BsR, between BR and BsR, between 2R and 4R, or between 4R and SR. However, the percentage duration of 2R was significantly higher than that of SR ($t_{24} = 2.15$, $P = 0.042$).

Both bark-scratching actions (2Sc and SSc) were observed on Japanese larch trees, and the damaged bark

Table 1. Percentages of visits between 2010 and 2015 in which Asiatic black bears (*Ursus thibetanus*) in Japan exhibited tree-marking or associated behaviors and the average (SD) percentage of time occupied by each behavior.

Type of tree-marking or associated behavior		% (no.) of visits in which behavior was displayed (N = 22)	Average % (SD) of time occupied by behavior during visits in which the behavior was displayed
Tree-rubbing actions			
4BsR	Quadrupedal standing position, body-side rubbing	59.1 (13)	55.3 (33.6)
2FR	Bipedal standing position, front rubbing	40.9 (9)	40.0 (25.5)
SFR	Sitting position, front rubbing	40.9 (9)	45.3 (17.2)
2BR	Bipedal standing position, back rubbing	59.1 (13)	54.1 (33.9)
SBR	Sitting position, back rubbing	4.5 (1)	10.2 (n/a) ^a
Tree-rubbing orientations			
BsR	Body-side rubbing	59.1 (13)	55.3 (33.6)
FR	Front rubbing	59.1 (13)	59.0 (21.9)
BR	Back rubbing	59.1 (13)	51.0 (35.2)
Tree-rubbing postures			
4R	Quadrupedal standing position, rubbing	59.1 (13)	55.3 (33.6)
2R	Bipedal standing position, rubbing	72.7 (16)	66.5 (32.2)
SR	Sitting position, rubbing	40.9 (9)	41.7 (20.8)
Debarking actions			
2Sc	Bipedal standing position, bark scratching	22.7 (5)	—
SSc	Sitting position, bark scratching	22.7 (5)	—
Associated behaviors			
St	Stomping	36.4 (8)	—

^aSitting position, back rubbing was observed only once.

was surrounded by a border entirely stripped of bark by repeated claw-scratching. No bark-biting behavior was observed. Of the 8 visits in which St was confirmed on the same Japanese larch tree (25 Jul to 8 Sep 2011), 6 were by the same male and 2 were by bears of unknown gender. Up to 13 stomps over the same spot were confirmed to have been made as the bears moved away from the target tree (Fig. S3, Supplemental material). We could not observe the urinating behavior.

In the analysis of the order of each action, 4BsR was shown to have the highest probability of being performed immediately after the start of the visit (57%; Fig. 1). Following 4BsR, leaving without St had the highest probability of occurring (48%), followed by 2BR (26%). Following 2BR, 2FR had the highest probability of occurring (38%). Actions that occurred after 2FR included leaving (45%) and 2Sc (36%), and St behavior had a high probability of occurring after 2Sc (60%). The probability of 2BR occurring after 2FR was 0%.

Discussion

To our knowledge, this is the first study of the tree-marking behavior of Asiatic black bears. We found that

there were no significant differences in the frequencies or durations of FR and either BR or BsR and of BR and BsR during tree-marking visits, but that a bipedal standing (2R) posture occurred in a high percentage of visits and had a high percentage duration. Other studies have shown that brown bears and American black bears are most likely to perform BR on 2 legs (2BR), rather than BsR while standing on 4 feet (4BsR) or rubbing while standing on 2 legs facing the tree (2FR; Clapham et al. 2014, Taylor et al. 2015). Notably, brown bears may perform scent-marking while sitting and rubbing their backs against the tree in order to mark it with their anal glands and sebaceous glands located on the back (Rosell et al. 2011, Tomiyasu et al. 2017). On the other hand, in a study of Andean bears (Filipczyková et al. 2016), the percentage of visits in which FR and BR were performed was about the same, similar to our findings with Asiatic black bears. In the Andean bear study, however, a lower percentage of 4BsR occurred. In any case, to date few analyses have been conducted on Asiatic black bear glandular secretions and their role in tree-marking.

We also examined the order in which tree-marking actions occurred to determine whether it is related to the ease of transition between actions or to the transition preference; however, it might not be explained by the ease

black bears (Ziegltrum 2009), and Asiatic black bears (Kobashikawa and Koike 2016). However, although both bark-scratching and cambial feeding are performed on the bark of conifers, it is unknown whether these behaviors are related to each other, and no cambial feeding was observed in this study. More research is needed to address these questions.

In summary, we studied the characteristics of the marking behavior of Asiatic black bears at rub trees and found possible differences from other bears in the body orientation tendency when rubbing against a tree and the order of marking behaviors, although the number of bears observed here was limited. To understand the unique characteristics of communication by scent-marking in Asiatic black bears, we need to study a much larger sample of bears, with behavioral analysis from a wider area and specific data on the presence, position, and composition of sebaceous glands and the relationship between them.

Acknowledgments

We would like to thank the members of Okutama Black Bear Research Group for supporting the field survey. We also thank Dr. Koji Yamazaki for providing information on bears and encouraging our research. We thank the Okutama Town Office for permission to set camera traps. This work was partially supported by JSPS KAKENHI (grant numbers JP16H04939, JP16H04932, and JP17H05971) and the Institute of Global Innovation Research at the Tokyo University of Agriculture and Technology. We thank the Editor in Chief Dr. Jon Swenson, the Associate Editor, and 3 anonymous reviewers for their valuable suggestions and comments on the earlier versions of this paper.

Literature cited

- AL TOMARE, D., AND D. LORIS. 2018. "Package 'Channel Attribution'." <https://cran.r-project.org/web/packages/ChannelAttribution/ChannelAttribution.pdf>. Accessed 27 Apr 2020.
- BURST, T.L., AND M.R. PELTON. 1983. Black bear mark trees in the Smoky Mountains. *Bears: Their Biology and Management* 5:45–53.
- CHOO, Y.R., E.P. KUDAVIDANAGE, T.R. AMARASINGHE, T. NIMALRATHNA, M.A.H. CHUA, AND E.L. WEBB. 2020. Best practices for reporting individual identification using camera trap photographs. *Global Ecology and Conservation* 24:e01294. <https://doi.org/10.1016/j.gecco.2020.e01294>.
- CLAPHAM, M., O.T. NEVIN, A.D. RAMSEY, AND F. ROSELL. 2012. A hypothetico-deductive approach to assessing the social function of chemical signalling in a non-territorial solitary carnivore. *PLoS One* 7:e35404. <https://doi.org/10.1371/journal.pone.0035404>.
- _____, _____, _____, AND _____. 2013. The function of strategic tree selectivity in the chemical signalling of brown bears. *Animal Behaviour* 85:1351–1357. <https://doi.org/10.1016/j.anbehav.2013.03.026>.
- _____, _____, _____, AND _____. 2014. Scent-marking investment and motor patterns are affected by the age and sex of wild brown bears. *Animal Behaviour* 94:107–116. <https://doi.org/10.1016/j.anbehav.2014.05.017>.
- FILIPCZYKOVÁ, E., I.M.A. HEITKONIG, A. CASTELLANOS, W. HANSTON, AND S.M.J.G. STEYAERT. 2016. Marking behavior of Andean bears in an Ecuadorian cloud forest: A pilot study. *Ursus* 27:122–128. <https://doi.org/10.2192/URSUS-D-16-00002.1>.
- GREEN, G.I., AND D.J. MATTSON. 2003. Tree rubbing by Yellowstone grizzly bears *Ursus arctos*. *Wildlife Biology* 9:1–9.
- JAPAN METEOROLOGICAL AGENCY. 2018. Past weather data research. Japan Meteorological Agency, Tokyo. [In Japanese.] www.jma.go.jp. Accessed 5 May 2018.
- JOJOLA, S.M., F. ROSELL, I. WARRINGTON, J.E. SWENSON, AND A. ZEDROSSER. 2012. Subadult brown bears (*Ursus arctos*) discriminate between unfamiliar adult male and female anal gland secretion. *Mammalian Biology* 77:363–368.
- KARAMANLIDIS, A.A., D. YOULATOS, S. SGARDELIS, AND Z. SCOURAS. 2007. Using sign at power poles to document presence of bears in Greece. *Ursus* 18:54–61. [https://doi.org/10.2192/1537-6176\(2007\)18\[54:USAPPT\]2.0.CO;2](https://doi.org/10.2192/1537-6176(2007)18[54:USAPPT]2.0.CO;2).
- KENDALL, C.K., J.B. STETZ, J. BOULANGER, A.C. MACLEOD, D. PAETKAU, AND G.C. WHITE. 2009. Demography and genetic structure of a recovering grizzly bear population. *Journal of Wildlife Management* 73:3–17. <https://doi.org/10.2193/2008-330>.
- KLEINER, J.D., R.C.V. HORN, J.E. SWENSON, AND S.M.J.G. STEYAERT. 2018. Rub-tree selection by Andean bears in the Peruvian dry forest. *Ursus* 29:58–66. <https://doi.org/10.2192/URSUS-D-17-00012.1>.
- KOBASHIKAWA, S., AND S. KOIKE. 2016. Spatiotemporal factors affecting bark stripping of conifer trees by Asiatic black bears (*Ursus thibetanus*) in Japan. *Forest Ecology and Management* 380:100–106. <https://doi.org/10.1016/j.foreco.2016.08.042>.
- KOIKE, S., H. MORIMOTO, Y. GOTO, C. KOZAKAI, AND K. YAMAZAKI. 2008. Frugivory of carnivores and seed dispersal of fleshy fruits in cool-temperate deciduous forests. *Journal of Forest Research* 13:215–222. <https://doi.org/10.1007/s10310-008-0069-5>.
- LATHAM, E., J.B. STETZ, I. SERYODKIN, D. MIQUELLE, AND M.L. GIBEAU. 2012. Non-invasive genetic sampling of brown bears and Asiatic black bears in the Russian Far East: A pilot study. *Ursus* 23:145–158.
- NIE, Y., R.R. SWAISGOOD, Z. ZHANG, Y. HU, Y. MA, AND F. WEI. 2012. Giant panda scent-marking strategies in the wild:

- Role of season, sex and marking surface. *Animal Behaviour* 84:39–44.
- OGAWA, Y., K. TOCHIGI, T. NAGANUMA, B.S. DEWI, AND S. KOIKE. 2020. Tree rubbing by Asian black bears (*Ursus thibetanus*) in conifer plantations in Okutama Mountain in Japan. *Animal Biology* 70:351–358. <https://doi.org/10.1163/15707563-bja10006>.
- R CORE TEAM. 2019. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>. Accessed 24 Mar 2020.
- ROSELL, F., S.M. JOJOLA, K. INGDAL, B.A. LASSEN, J.E. SWENSON, J.M. ARNEMO, AND A. ZEDROSSER. 2011. Brown bears possess anal sacs and secretions may code for sex. *Journal of Zoology* 283:143–152. <https://doi.org/10.1111/j.1469-7998.2010.00754.x>.
- SATO, Y., C. KAMIISHI, T. TOKAJI, M. MORI, S. KOIZUMI, K. KOBAYASHI, T. ITOH, W. SONOHARA, M.B. TAKADA, AND T. URATA. 2014. Selection of rub trees by brown bears (*Ursus arctos*) in Hokkaido, Japan. *Acta Theriologica* 59:129–137. <https://doi.org/10.1007/s13364-013-0143-z>.
- SERGIEL, A., J. NAVES, P. KUJAWSKI, R. MAŚLAK, E. SERWA, D. RAMOS, A. FERNAÁNDEZ-GIL, E. REVILLA, T. ZWIJACZ-KOZICA, F. ZIEĘBA, J. PAINER, AND N. SELVA. 2017. Histological, chemical and behavioural evidence of pedal communication in brown bears. *Scientific Reports* 7:1052. <https://doi.org/10.1038/s41598-017-01136-1>.
- TATTONI, C., N. BRAGALANTI, C. GROFF, AND F. ROVERO. 2015. Patterns in the use of rub trees by the Eurasian brown bear. *Hystrix, the Italian Journal of Mammalogy* 26:118–124. <https://doi.org/10.4404/hystrix-26.2-11414>.
- TAYLOR, A.P., M.L. ALLEN, AND M.S. GUNTHER. 2015. Black bear marking behaviour at rub trees during the breeding season in northern California. *Behaviour* 152:1097–1111. <https://doi.org/10.1163/1568539X-00003270>.
- THIEURMEL, B. 2016. visNetwork: Network Visualization using 'vis.js' Library. Rpackageversion1.0.2. <https://CRAN.R-project.org/package=visNetwork>. Accessed 24 Mar 2020.
- TOMIYASU, J., Y. YANAGAWA, Y. SATO, M. SHIMOZURU, M. NAGANO, M. SASAKI, H. SAKAMOTO, N. MATSUMOTO, K. KOBAYASHI, M. KAYANO, S. HANEDA, AND M. MATSUI. 2017. Testosterone-related and seasonal changes in sebaceous glands in the back skin of adult male brown bears (*Ursus arctos*). *Canadian Journal of Zoology* 96:205–211. <https://doi.org/10.1139/cjz-2017-0028>.
- WHITE, A.M., R.R. SWAISGOOD, AND H. ZHANG. 2002. The highs and lows of chemical communication in giant pandas (*Ailuropoda melanoleuca*): Effect of scent deposition height on signal discrimination. *Behavioral Ecology and Sociobiology* 51:519–529.
- ZIEGLTRUM, G.J. 2009. Evaluation of the black bear supplemental feeding program in western Washington, USA. Dissertation, Technical University of Munich, Germany.
- ZYŚK-GORCZYŃSKA, E., Z. JAKUBIEC, B. WERTZ, AND A. WUCZYŃSKI. 2016. Long-term study of damage to trees by brown bears *Ursus arctos* in Poland: Increasing trends with insignificant effects on forest management. *Forest Ecology and Management* 366:53–64. <https://doi.org/10.1016/j.foreco.2016.02.007>.

Received: August 26, 2020

Accepted: February 10, 2021

Associate Editor: V. Penteriani

Supplemental material

Fig. S1. Map of the study area and survey site in Okutama, Tokyo, Japan, where we used automatic cameras to record tree-marking and associated behavior of Asiatic black bears (*Ursus thibetanus*) between 2010 and 2015.

Fig. S2. Classification of tree-marking actions in Asiatic black bears (*Ursus thibetanus*).

Fig. S3. Sample photographs of steps 1–13 of the stomping behavior by an Asiatic black bear (*Ursus thibetanus*) in Japan that occurred after tree-rubbing behavior on 25 July 2011, at 1842 hours.